



# Chem 108:

## Lab

### Week 14


Sign in  
Note your Group #  
Pick up Papers  
and Handout

Week 14	Lab Notes:
	<div></div> <p>Lab 2341.14:</p> <p>Discussion, Experiments &amp; Graded Assignments:</p> <p>Week #14 Powerpoint <a href="#">.html</a>, <a href="#">.pptx</a>, Print: <a href="#">.pdf</a> (6 slides per page)</p> <div></div> <p>Experimentation:</p> <p>Doing:</p> <p><i>Gas Stoichiometry, Lab Manual</i> pp.53-60; <a href="#">Procedure</a> pp.53-57; <a href="#">Report Form</a> pp.58-60 Part I pg. 58 + Data pg. 59 DUE Today; Completed pp.58-59, DUE 3-Dec</p> <ul style="list-style-type: none"><li>• (GQ) Viewing: <a href="#">Acids-Bases pH Guiding Questions</a></li><li>• <a href="#">Fluid Exchange/ Fluid Exchange Form</a> DUE Today</li><li>• Household Acids &amp; Bases / pH: <a href="#">Experiment &amp; Report Form</a> DUE Today <a href="#">Post Lab on-line Acid-Base Descriptions</a> DUE Today</li></ul> <p><i>Acid-Base Titration: Laboratory Manual</i> <a href="#">Acid &amp; Base Titration</a> (pp. 91-93) Part II: <a href="#">Individual Unknown</a> (pg. 95 <a href="#">Completed form</a>) DUE Today</p>

# Gas Stoichiometry

<http://chemconnections.org/general/chem108/Magnesium-Zinc-wo.1.mov>

## *Experimentally Determining Moles of Hydrogen*



## *Using Partial Pressures the Ideal Gas Law & Stoichiometry*

*Dr. Ron Rusay*



Except where otherwise [noted](#), content on this site is licensed under a [Creative Commons Attribution 4.0 International license](#).

- Refer to the Procedure section pp. 54-56. The following slides correspond to the instructions in the procedure.

Chem 108/ Dr. Rusay

#### Equipment

100 mL eudiometer  
buret clamp  
red alcohol thermometer  
ruler  
ring stand  
large beaker (at least 400 mL)  
wash bottle w/ deionized water

#### Procedure

Refer to the on-line movie and the on-line notes for today's class, and then complete Part I of the Report Form. After completing Part I, obtain a metal sample envelope from Dr. R. Record its number and the mass of magnesium in the report form. Make a cage around the piece of magnesium using fine copper wire. First fold the ribbon several times to make it as compact as possible. NOTE: The cage must be tight enough so that the metal cannot fall out as it reacts and loses size. If too much wire is used and the cage is too tight, the reaction may be very slow. Leave a tail of copper wire about 10 cm long. Pour approximately 20 mL of dilute (6 M) hydrochloric acid into a clean 100 mL eudiometer. This does not need to be measured accurately nor does the exact volume need to be known. Carefully and slowly fill the rest of the eudiometer with deionized water so as to avoid mixing of the water and the acid. Insert the magnesium sample in the eudiometer so that it is ~ 10 cm from the stopper (when it is upside down) and fix its position by placing the copper wire tail against the wall of the eudiometer pressing against a one-hole rubber stopper as illustrated in the presentation. When inserting the rubber stopper, let the excess water come out through the hole. Make sure no air is trapped in the tube as it will later be measured as hydrogen gas causing error. Cover the hole in the stopper with your finger and invert the eudiometer in a large beaker partly filled with water and clamp it to a ring stand using a buret clamp. The acid solution, being denser than the water, mixes slowly and concentrates down the eudiometer until it reacts with the metal producing hydrogen gas.










Figure 1—Eudiometer


What is wrong with this set up?








**Pressure on the inside  
of the eudiometer  
must equal the pressure  
outside the eudiometer!**



*Figure 2*


- Refer to the Gas Stoichiometry Report Form, pg. 58-59
- Experimental data is to be obtained for the reaction of a known mass of magnesium metal:



- The volume of hydrogen, pressure and temperature determined and recorded.
- Moles of hydrogen is calculated using Ideal Gas Law calculations, then calculating mass of the starting magnesium from the number of moles of hydrogen.



# Background Ideal Gas Law



$$PV = nRT$$

- $R$  = “proportionality” constant  
=  $0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$
- $P$  = pressure of gas in atm
- $V$  = volume of gas in liters
- $n$  = moles of gas
- $T$  = temperature of gas in Kelvin


# *Standard Conditions*

## *Temperature, Pressure & Moles*


- “STP”

- For 1 mole of a gas at STP:
- $P = 1$  atmosphere
- $T = 0^{\circ}\text{C}$  (273.15 K)
- The molar volume of an ideal gas is **22.42** liters at STP

Isobaric process: pressure constant  
 Isochoric process: volume constant  
 Isothermal process: temperature constant



# *Ideal Gas Law Simulator*





<http://ch301.cm.utexas.edu/simulations/gas-laws/GasLawSimulator.swf>

# *Hydrogen & the Ideal Gas Law*

$$n_{\text{H}_2(\text{g})} = PV / RT$$

- $n$  = moles  $\text{H}_2(\text{g})$
- $P_{\text{H}_2(\text{g})}$  = pressure of  $\text{H}_2(\text{g})$  in atm (mm Hg  $\rightarrow$  atm)
- $V$  = experimental volume (mL  $\rightarrow$  L)
- $T$  = experimental temperature ( $^{\circ}\text{C} \rightarrow \text{K}$ )




# *Total Pressure:*

## *Sum of the Partial Pressures*

- For a mixture of gases, the total pressure is the sum of the pressures of each gas in the mixture.

$$P_{\text{Total}} = P_1 + P_2 + P_3 + \dots$$

$$P_{\text{Total}} \propto n_{\text{Total}}$$



$$n_{\text{Total}} = n_1 + n_2 + n_3 + \dots$$


$$\bullet P_{H_2(g)} = P_{\text{Total (barometric)}} - P_{H_2O(g)} [TABLE] - P_{HCl(g)}$$

$$P_{HCl(g)} =$$

$$HCl \text{ Height (mm)} \div 12.95$$


---

*Density Hg is  
12.95 times >  
density HCl(aq)*



$$P_{HCl(g)} =$$

$$HCl \text{ Height (mm)} \times 0.0772$$


---

*Density Hg is  
12.95 times >  
density HCl(aq)*

*0.772 mm Hg/cm of acid solution*





# *Ideal Gas Law:*


## *Moles / Avogadro's Law*

$$n_{\text{H}_2(\text{g})} = \frac{PV}{RT}$$

- $n$  = moles  $\text{H}_2(\text{g})$
- $P_{\text{H}_2(\text{g})}$  = pressure of  $\text{H}_2(\text{g})$  in atm (mm Hg  $\rightarrow$  atm)
- $P_{\text{H}_2(\text{g})} = P_{\text{Total}}$  (barometric) -  $P_{\text{H}_2\text{O}(\text{g})}$  [TABLE] -  $P_{\text{HCl}(\text{g})}$
- $V$  = experimental volume (mL  $\rightarrow$  L)
- $T$  = experimental temperature ( $^{\circ}\text{C} \rightarrow \text{K}$ )
- $R = 0.082057338 \text{ L atm K}^{-1} \text{ mol}^{-1}$  (constant)



- Refer to Report Form Part I: (Example uses Zinc.)




Mole Calculations:

- Stoichiometry Calculation
- Ideal Gas Law Calculations
- Comparison (% Error)

# *Stoichiometry*

## *Moles Hydrogen / Mass of Zinc* (Part I: Zinc Calculation)




$$\text{mol H}_2\text{(g)} = \text{mol Zn(s)}$$

$$\text{mass (g) Zn(s)} = \text{mol Zn(s)} \times \text{Molar Mass Zn(s)}$$

# Zinc Example Calculation

- Complete Report Form pg. 58 Part I:



Mole Calculations:

- Stoichiometry Calculation
- Ideal Gas Law Calculations
- Comparison (% Error)

## Report Form – Gas Stoichiometry

### Part I – Sample Data for Mass of Zinc


Chemical Reaction		
DATA COLLECTED		
Volume of hydrogen collected*	81.5 mL	L
Temperature of hydrogen*	22.0 °C	K
Barometric pressure*	29.98 in Hg	mm Hg
Height of solution in eudiometer from benchtop	19.2 cm	
Height of solution in beaker from benchtop	10.0 cm	
CALCULATIONS AND RESULTS		
Difference in liquid levels of solution in eudiometer and beaker*		
Aqueous vapor pressure at temperature of hydrogen	mm Hg	
Pressure caused by acid column:*( (Difference in cm)*(0.772 mm Hg/cm)	mm Hg	
Pressure of hydrogen alone*	mm Hg	atm
Moles of hydrogen*	moles	
Moles of zinc*	moles	
Mass of zinc (calculated)*	g	

Show the calculations for each of the entries in the Data Table marked with \* on the calculations page.

**Question:** If the mass of zinc used was 0.21 g, what is the percent error for your calculated mass of zinc? Show your work below.

# *Moles : Ideal Gas Law*

(Part I: Zinc Calculation Example)




$$n_{\text{H}_2\text{(g)}} = PV / RT$$

- **n = moles  $\text{H}_2\text{(g)}$**
- **P  $\text{H}_2\text{(g)}$  = pressure of  $\text{H}_2\text{(g)}$  in atm (mm Hg  $\rightarrow$  atm)**
- **P  $\text{H}_2\text{(g)}$  = P Total (barometric) - P  $\text{H}_2\text{O}$  (g) [TABLE] - P HCl (g)**
- **V = experimental volume (mL  $\rightarrow$  L)**
- **T = experimental temperature ( $^{\circ}\text{C} \rightarrow \text{K}$ )**

$$\mathbf{R = 0.082057338 \text{ L atm K}^{-1} \text{ mol}^{-1}}$$

# Moles : Ideal Gas Law

## (Part I: Zinc Calculation Example)



$$n_{\text{H}_2(\text{g})} = PV / RT$$

**V** = experimental volume  
(mL → L)

$$R = 0.082057338 \text{ L atm K}^{-1} \text{ mol}^{-1}$$

### Report Form – Gas Stoichiometry

#### Part I – Sample Data for Mass of Zinc


Chemical Reaction		
DATA COLLECTED		
Volume of hydrogen collected*	81.5 mL	
Temperature of hydrogen*	22.0 °C	
Barometric pressure*	29.98 in Hg	mm Hg
Height of solution in eudiometer from benchtop	19.2 cm	
Height of solution in beaker from benchtop	10.0 cm	
CALCULATIONS AND RESULTS		
Difference in liquid levels of solution in eudiometer and beaker*		
Aqueous vapor pressure at temperature of hydrogen	mm Hg	
Pressure caused by acid column:*(Difference in cm)*(0.772 mm Hg/cm)	mm Hg	
Pressure of hydrogen alone*	mm Hg	atm
Moles of hydrogen*	moles	
Moles of zinc*	moles	
Mass of zinc (calculated)*	g	

Show the calculations for each of the entries in the Data Table marked with \* on the calculations page.

**Question:** If the mass of zinc used was 0.21 g, what is the percent error for your calculated mass of zinc? Show your work below.

# Moles : Ideal Gas Law

## (Part I: Zinc Calculation Example)



$$n_{\text{H}_2\text{(g)}} = PV / RT$$

**V** = experimental volume  
(mL → L)

**T** = experimental temperature  
(°C → K)

$$R = 0.082057338 \text{ L atm K}^{-1} \text{ mol}^{-1}$$

### Report Form – Gas Stoichiometry

#### Part I – Sample Data for Mass of Zinc

Chemical Reaction			
DATA COLLECTED			
Volume of hydrogen collected*	81.5 mL		
Temperature of hydrogen*	22.0 °C		
Barometric pressure*	29.98 in Hg		mm Hg
Height of solution in eudiometer from benchtop			19.2 cm
Height of solution in beaker from benchtop			10.0 cm
CALCULATIONS AND RESULTS			
Difference in liquid levels of solution in eudiometer and beaker*			
Aqueous vapor pressure at temperature of hydrogen			mm Hg
Pressure caused by acid column:*( (Difference in cm)*(0.772 mm Hg/cm)	mm Hg		
Pressure of hydrogen alone*	mm Hg		atm
Moles of hydrogen*			moles
Moles of zinc*			moles
Mass of zinc (calculated)*			g


Show the calculations for each of the entries in the Data Table marked with \* on the calculations page.

**Question:** If the mass of zinc used was 0.21 g, what is the percent error for your calculated mass of zinc? Show your work below.



# Moles : Ideal Gas Law

## (Part I: Zinc Calculation Example)



$$n_{\text{H}_2(\text{g})} = PV / RT$$

**V** = experimental volume  
(mL → L)

**T** = experimental temperature  
(°C → K)

**P**  $\text{H}_2(\text{g})$  = pressure of  $\text{H}_2(\text{g})$  in  
atm (mm Hg → atm)

**P**  $\text{H}_2(\text{g})$  = **P** Total (barometric) - **P**  
 $\text{H}_2\text{O}(\text{g})$  [TABLE] - **P**  $\text{HCl}(\text{g})$

$$R = 0.082057338 \text{ L atm K}^{-1} \text{ mol}^{-1}$$

### Report Form – Gas Stoichiometry

#### Part I – Sample Data for Mass of Zinc


Chemical Reaction			
DATA COLLECTED			
Volume of hydrogen collected*	81.5 mL		L
Temperature of hydrogen*	22.0 °C		K
Barometric pressure*	29.98 in Hg		mm Hg
Height of solution in eudiometer from benchtop			19.2 cm
Height of solution in beaker from benchtop			10.0 cm
CALCULATIONS AND RESULTS			
Difference in liquid levels of solution in eudiometer and beaker*			
Aqueous vapor pressure at temperature of hydrogen	mm Hg		
Pressure caused by acid column:*(Difference in cm)*(0.772 mm Hg/cm)	mm Hg		
Pressure of hydrogen alone*	mm Hg		atm
Moles of hydrogen*	moles		
Moles of zinc*	moles		
Mass of zinc (calculated)*	g		

Show the calculations for each of the entries in the Data Table marked with \* on the calculations page.

**Question:** If the mass of zinc used was 0.21 g, what is the percent error for your calculated mass of zinc? Show your work below.

# Moles : Ideal Gas Law

Part I: Hydrogen Calculation, (*Refer to Form's Data*)




$$n_{\text{H}_2\text{(g)}} = \frac{PV}{RT}$$

- $n$  = moles  $\text{H}_2\text{(g)}$
- $P_{\text{H}_2\text{(g)}}$  = pressure of  $\text{H}_2\text{(g)}$  in atm (mm Hg  $\rightarrow$  atm)
- $P_{\text{H}_2\text{(g)}} = 29.98$  inches Hg (barometric) - 19.8 mm Hg  $\text{H}_2\text{O (g)}$  [TABLE]  
-  $P_{\text{HCl (g)}}$

$P_{\text{HCl (g)}}$

$$R = 0.082057338 \text{ L atm K}^{-1} \text{ mol}^{-1}$$



$$\bullet P_{H_2(g)} = P_{\text{Total (barometric)}} - P_{H_2O(g)} [TABLE] - P_{HCl(g)}$$

$$P_{HCl(g)} =$$

$$19.2 \text{ cm Hg} - 10.0 \text{ cm Hg} = 9.2 \text{ cm Hg}$$


$$HCl \text{ Height (mm)} \div 12.95$$

$$= 7.1 \text{ mm Hg}$$

---

*Density Hg is  
12.95 times >  
density HCl(aq)*

$$P_{HCl(g)}$$



*0.772 mm Hg/cm of acid solution*

$$P_{HCl(g)} =$$

$$19.2 \text{ cm Hg} - 10.0 \text{ cm Hg} = 9.2 \text{ cm Hg}$$


$$HCl \text{ Height (mm)} \times 0.0772$$

$$= 7.1 \text{ mm Hg}$$

---


*Density Hg is  
12.95 times >  
density HCl(aq)*

$$\begin{aligned}
 P_{\text{H}_2(\text{g})} &= 761.5 \text{ mm Hg (barometric)} \\
 &- 19.8 \text{ mm Hg H}_2\text{O (g)} - 7.1 \text{ mm Hg HCl (g)} \\
 &= 734.6 \text{ mm Hg} \\
 &= 734.6 \text{ mm Hg} / 760.0 \text{ mm Hg} / 1.000 \text{ atm} \\
 &= 0.9666 \text{ atm}
 \end{aligned}$$



# *Moles : Ideal Gas Law*

(Part I: Hydrogen Calculation)



$$n_{\text{H}_2\text{(g)}} = \frac{PV}{RT}$$

- $n = \text{moles H}_2\text{(g)}$
- $P_{\text{H}_2\text{(g)}} = 0.9666 \text{ atm}$
- $V = 0.0815 \text{ L}$
- $T = 295.1 \text{ K}$


$$R = 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$$

$$n_{\text{H}_2\text{(g)}} = 0.00325 \text{ moles H}_2\text{(g)} = 0.00325 \text{ moles Zn(s)}$$

# *% Error*

## *Theoretical Mass Zinc vs. Experimental*

(Part I: Calculation)



$$\text{mass (g) Zn(s)} = \text{mol Zn(s)} \times \text{Molar Mass Zn(s)}$$

$$= 0.00325 \text{ moles Zn(s)} \times 65.37 \text{ g/mol Zn(s)}$$

$$\% \text{ Error} = \frac{\text{experimental grams Zn(s)} - \text{theoretical grams Zn(s)}}{\text{theoretical grams Zn(s)}} \times 100$$

$$= \frac{0.213 \text{ g} - 0.21 \text{ g}}{0.21 \text{ g}} \times 100$$

$$= 1.4 \%$$

Bring completed Report Forms to Dr. R. to get Mg(s) sample(s).


# *Molar Mass of any Gas*

*(Hydrogen for example)*

- $PV = nRT$
- $n = \text{g of gas} / \text{MM}_{\text{gas}}$  [ $\text{MM}_{\text{gas}} = \text{g/mol}$ ]
- $PV = (\text{g of gas} / \text{MM}_{\text{gas}})RT$
- $\text{MM}_{\text{gas}} = \text{g of gas} / V (RT/P)$


## *Density of gas*

- $\text{MM}_{\text{gas}} = \text{g of gas} / V (RT/P)$
- $\text{MM}_{\text{gas}} = \text{density of gas} (RT/P)$





# (Part II) Magnesium



Mole Calculations:

- Stoichiometry Calculation
- Ideal Gas Law Calculations
- Comparison (% Error)

Get equipment from stockroom and complete data acquisition for Part II.

Have individual Report Forms checked before leaving lab today.

Name: \_\_\_\_\_

Section: \_\_\_\_\_

## Part II - Mass of Magnesium

Chemical Reaction		
<b>DATA COLLECTED</b>		
Unknown number		
Volume of hydrogen collected*	mL	L
Temperature of hydrogen*	°C	K
Barometric pressure*	inches Hg	mm Hg
Height of solution in eudiometer from benchtop	cm	
Height of solution in beaker from benchtop	cm	
<b>CALCULATIONS AND RESULTS</b>		
Difference in liquid levels of solution in eudiometer and beaker*	cm Acid Solution	
Aqueous vapor pressure at temperature of hydrogen	mm Hg	
Pressure caused by acid column:* (Difference in cm) * (0.772 mmHg/cm)	mm Hg	
Pressure of hydrogen alone*	mm Hg	atm
Moles of hydrogen*	moles	
Moles of magnesium*	moles	
Mass of magnesium*	g	

Show the calculations for each of the entries in the Data Table marked with \* on the calculations page.