


Chem 108: Lab Week 14



Sign in
Note your Group #
Pick up Papers
and Handout



Gas Stoichiometry

<http://chemconnections.org/general/chem108/Magnesium-Zinc-wo.1.mov>

Experimentally Determining Moles of Hydrogen



Using Partial Pressures the Ideal Gas Law & Stoichiometry

Dr. Ron Ruxay



Except where otherwise noted, content on this site is licensed under a Creative Commons Attribution 4.0 International License.

- Refer to the Procedure section pp. 54-56. The following slides correspond to the instructions in the procedure.

Chem 108/Dr. Ruxay

Equipment

100 mL eudiometer
beaker clamp
red alcohol thermometer
ruler
ring stand
large beaker (at least 400 mL)
wash bottle w/ deionized water

Procedure

Refer to the on-line movie and the on-line notes for today's class, and then complete Part 1 of the Report Form. After completing Part 1, obtain a metal sample envelope from Dr. R. Record its number and the mass of magnesium in the report form. Make a cage around the piece of magnesium using fine copper wire. First fold the ribbon several times to make it as compact as possible. NOTE: The cage must be tight enough so that the metal cannot fall out as it reacts and loses size. If no mesh wire is used and the cage is too tight, the reaction may be very slow. Leave a tail of copper wire about 10 cm long. Place approximately 30 mL of dilute (6 M) hydrochloric acid into a clean 100 mL eudiometer. This does not need to be measured accurately nor does the exact volume need to be known. Carefully and slowly fill the rest of the eudiometer with deionized water so as to avoid mixing of the water and the acid. Insert the magnesium sample in the eudiometer so that it is ~10 cm from the copper (when it is upside down) and fix its position by placing the copper wire tail against the wall of the eudiometer pressing against a cone-hole rubber stopper as illustrated in the presentation. When inserting the rubber stopper, let the excess water come out through the hole. Make sure no air is trapped in the tube as it will later be measured as hydrogen gas causing error. Cover the hole in the stopper with your finger and invert the eudiometer in a large beaker partly filled with water and clamp it to a ring stand using a beaker clamp. The acid solution, being denser than the water, mixes slowly and concentrates down the eudiometer until it reaches with the metal producing hydrogen gas.








Figure 1-Eudiometer

What is wrong with this set up?





- Refer to the Gas Stoichiometry Report Form, pg. 58-59
- Experimental data is to be obtained for the reaction of a known mass of magnesium metal:

$$\text{Mg(s)} + 2\text{HCl(aq)} \rightarrow \text{MgCl}_2\text{(aq)} + \text{H}_2\text{(g)}$$
- The volume of hydrogen, pressure and temperature determined and recorded.
- Moles of hydrogen is calculated using Ideal Gas Law calculations, then calculating mass of the starting magnesium from the number of moles of hydrogen.

Background
Ideal Gas Law


$PV = nRT$

- $R = \text{"proportionality" constant} = 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$
- $P = \text{pressure of gas in atm}$
- $V = \text{volume of gas in liters}$
- $n = \text{moles of gas}$
- $T = \text{temperature of gas in Kelvin}$


Standard Conditions Temperature, Pressure & Moles

• "STP"

- For 1 mole of a gas at STP:
- $P = 1 \text{ atmosphere}$
- $T = 0^\circ\text{C} \text{ (273.15 K)}$
- The molar volume of an ideal gas is **22.42** liters at STP



Ideal Gas Law Simulator





<http://ch301.cm.utexas.edu/simulations/gas-laws/GasLawSimulator.swf>

Hydrogen & the Ideal Gas Law

$$n_{H_2(g)} = PV / RT$$

- n = moles $H_2(g)$
- $P_{H_2(g)}$ = pressure of $H_2(g)$ in atm (mm Hg $\rightarrow$ atm)
- V = experimental volume (mL $\rightarrow$ L)
- T = experimental temperature ($^{\circ}\text{C} \rightarrow$ K)




Total Pressure: Sum of the Partial Pressures

- For a mixture of gases, the total pressure is the sum of the pressures of each gas in the mixture.

$$P_{\text{Total}} = P_1 + P_2 + P_3 + \dots$$

$$P_{\text{Total}} \propto n_{\text{Total}}$$




$$n_{\text{Total}} = n_1 + n_2 + n_3 + \dots$$

$$P_{H_2(g)} = P_{\text{Total (barometric)}} - P_{H_2O(g)} [\text{TABLE}] - P_{HCl(g)}$$

$$P_{HCl(g)} =$$

$$\frac{\text{HCl Height (mm)}}{12.95}$$

Density Hg is
12.95 times >
density HCl(aq)



$$P_{HCl(g)} =$$

$$\frac{\text{HCl Height (mm)}}{0.0772}$$



Density Hg is
12.95 times >
density HCl(aq)

0.772 mm Hg/cm of acid solution


Ideal Gas Law: Moles / Avogadro's Law

$$n_{H_2(g)} = PV / RT$$

- n = moles $H_2(g)$
- $P_{H_2(g)}$ = pressure of $H_2(g)$ in atm (mm Hg $\rightarrow$ atm)
- $P_{H_2(g)} = P_{\text{Total (barometric)}} - P_{H_2O(g)} [\text{TABLE}] - P_{HCl(g)}$
- V = experimental volume (mL $\rightarrow$ L)
- T = experimental temperature ($^{\circ}\text{C} \rightarrow$ K)
- $R = 0.082057338 \text{ L atm K}^{-1} \text{ mol}^{-1}$ (constant)



- Refer to Report Form Part I: (Example uses Zinc.)




Mole Calculations:

- Stoichiometry Calculation
- Ideal Gas Law Calculations
- Comparison (% Error)

Stoichiometry

Moles Hydrogen / Mass of Zinc
(Part I: Zinc Calculation)



$$\text{mol } H_{2(g)} = \text{mol } Zn(s)$$

$$\text{mass (g) } Zn(s) = \text{mol } Zn(s) \times \text{Molar Mass } Zn(s)$$

Zinc Example Calculation

- Complete Report Form pg. 58 Part I:
- $$\text{Zn(s)} + 2\text{HCl(aq)} \rightarrow \text{ZnCl}_2\text{(aq)} + \text{H}_2\text{(g)}$$

Mole Calculations:

- Stoichiometry Calculation
- Ideal Gas Law Calculations
- Comparison (% Error)

Report Form - Gas Stoichiometry


Part I - Sample Data for Mass of Zinc

Chemical Reaction	
DATA COLLECTED	
Volume of hydrogen collected*	81.0 mL L
Temperature of hydrogen*	22.0 °C °C
Barometric pressure*	29.98 in Hg mm Hg
Height of solution in eudiometer from benchtop	19.2 cm
Height of solution in beaker from benchtop	10.0 cm
CALCULATIONS AND RESULTS	
Difference in liquid levels of solution in eudiometer and beaker*	mm Hg
Aqueous vapor pressure at temperature of hydrogen	mm Hg
Pressure caused by acid solution*	mm Hg
(Difference in level/10.0 mm Hg)	mm Hg
Pressure of hydrogen gas*	mm Hg
Pressure of hydrogen gas*	atm
Moles of hydrogen*	moles
Moles of zinc*	moles
Mass of zinc (calculated)*	g

Show the calculations for each of the entries in the Data Table marked with * on the calculation page.

Question: If the mass of zinc used was 0.21 g, what is the percent error for your calculated mass of zinc? Show your work below.

Moles : Ideal Gas Law (Part I: Zinc Calculation Example)




$$n_{\text{H}_2(\text{g})} = PV / RT$$

- n** = moles $\text{H}_2(\text{g})$
- P $\text{H}_2(\text{g})$** = pressure of $\text{H}_2(\text{g})$ in atm (mm Hg $\rightarrow$ atm)
- P $\text{H}_2(\text{g})$** = P Total (barometric) - P $\text{H}_2\text{O}(\text{g})$ [TABLE] - P HCl (g)
- V** = experimental volume (mL $\rightarrow$ L)
- T** = experimental temperature ($^{\circ}\text{C} \rightarrow \text{K}$)

$$R = 0.082057338 \text{ Latm K}^{-1} \text{ mol}^{-1}$$

Moles : Ideal Gas Law (Part I: Zinc Calculation Example)



$$n_{\text{H}_2(\text{g})} = PV / RT$$

V = experimental volume
(mL $\rightarrow$ L)

$$R = 0.082057338 \text{ Latm K}^{-1} \text{ mol}^{-1}$$

Report Form - Gas Stoichiometry


Part I - Sample Data for Mass of Zinc

Chemical Reaction	
DATA COLLECTED	
Volume of hydrogen collected*	81.0 mL L
Temperature of hydrogen*	22.0 °C °C
Barometric pressure*	29.98 in Hg mm Hg
Height of solution in eudiometer from benchtop	19.2 cm
Height of solution in beaker from benchtop	10.0 cm
CALCULATIONS AND RESULTS	
Difference in liquid levels of solution in eudiometer and beaker*	mm Hg
Aqueous vapor pressure at temperature of hydrogen	mm Hg
Pressure caused by acid solution*	mm Hg
(Difference in level/10.0 mm Hg)	mm Hg
Pressure of hydrogen gas*	mm Hg
Pressure of hydrogen gas*	atm
Moles of hydrogen*	moles
Moles of zinc*	moles
Mass of zinc (calculated)*	g

Show the calculations for each of the entries in the Data Table marked with * on the calculation page.

Question: If the mass of zinc used was 0.21 g, what is the percent error for your calculated mass of zinc? Show your work below.

Moles : Ideal Gas Law (Part I: Zinc Calculation Example)



$$n_{\text{H}_2(\text{g})} = PV / RT$$

V = experimental volume
(mL $\rightarrow$ L)

T = experimental temperature
($^{\circ}\text{C} \rightarrow \text{K}$)

$$R = 0.082057338 \text{ Latm K}^{-1} \text{ mol}^{-1}$$

Report Form - Gas Stoichiometry


Part I - Sample Data for Mass of Zinc

Chemical Reaction	
DATA COLLECTED	
Volume of hydrogen collected*	81.0 mL L
Temperature of hydrogen*	22.0 °C °C
Barometric pressure*	29.98 in Hg mm Hg
Height of solution in eudiometer from benchtop	19.2 cm
Height of solution in beaker from benchtop	10.0 cm
CALCULATIONS AND RESULTS	
Difference in liquid levels of solution in eudiometer and beaker*	mm Hg
Aqueous vapor pressure at temperature of hydrogen	mm Hg
Pressure caused by acid solution*	mm Hg
(Difference in level/10.0 mm Hg)	mm Hg
Pressure of hydrogen gas*	mm Hg
Pressure of hydrogen gas*	atm
Moles of hydrogen*	moles
Moles of zinc*	moles
Mass of zinc (calculated)*	g

Show the calculations for each of the entries in the Data Table marked with * on the calculation page.

Question: If the mass of zinc used was 0.21 g, what is the percent error for your calculated mass of zinc? Show your work below.

Moles : Ideal Gas Law (Part I: Zinc Calculation Example)



$$n_{\text{H}_2(\text{g})} = PV / RT$$

V = experimental volume
(mL $\rightarrow$ L)

T = experimental temperature
($^{\circ}\text{C} \rightarrow \text{K}$)

P $\text{H}_2(\text{g})$ = pressure of $\text{H}_2(\text{g})$ in
atm (mm Hg $\rightarrow$ atm)

P $\text{H}_2(\text{g})$ = P Total (barometric) - P
 $\text{H}_2\text{O}(\text{g})$ [TABLE] - P HCl (g)

$$R = 0.082057338 \text{ Latm K}^{-1} \text{ mol}^{-1}$$

Report Form - Gas Stoichiometry


Part I - Sample Data for Mass of Zinc

Chemical Reaction	
DATA COLLECTED	
Volume of hydrogen collected*	81.0 mL L
Temperature of hydrogen*	22.0 °C °C
Barometric pressure*	29.98 in Hg mm Hg
Height of solution in eudiometer from benchtop	19.2 cm
Height of solution in beaker from benchtop	10.0 cm
CALCULATIONS AND RESULTS	
Difference in liquid levels of solution in eudiometer and beaker*	mm Hg
Aqueous vapor pressure at temperature of hydrogen	mm Hg
Pressure caused by acid solution*	mm Hg
(Difference in level/10.0 mm Hg)	mm Hg
Pressure of hydrogen gas*	mm Hg
Pressure of hydrogen gas*	atm
Moles of hydrogen*	moles
Moles of zinc*	moles
Mass of zinc (calculated)*	g

Show the calculations for each of the entries in the Data Table marked with * on the calculation page.

Question: If the mass of zinc used was 0.21 g, what is the percent error for your calculated mass of zinc? Show your work below.

Moles : Ideal Gas Law Part I: Hydrogen Calculation, (Refer to Form's Data)




$$n_{\text{H}_2(\text{g})} = PV / RT$$

- n** = moles $\text{H}_2(\text{g})$
- P $\text{H}_2(\text{g})$** = pressure of $\text{H}_2(\text{g})$ in atm (mm Hg $\rightarrow$ atm)
- P $\text{H}_2(\text{g})$** = 29.98 inches Hg (barometric) - 19.8 mm Hg $\text{H}_2\text{O}(\text{g})$ [TABLE] - P HCl (g)


P HCl (g)

$$R = 0.082057338 \text{ Latm K}^{-1} \text{ mol}^{-1}$$



• $P_{H_2(g)} = P_{\text{Total (barometric)}} - P_{H_2O(g)} [\text{TABLE}] - P_{HCl(g)}$

$P_{HCl(g)} =$
 $19.2 \text{ cm Hg} - 10.0 \text{ cm Hg} = 9.2 \text{ mm Hg}$
 $HCl \text{ Height (mm)} \div 12.95$
 $= 7.1 \text{ mm Hg}$




$P_{HCl(g)} =$
 $19.2 \text{ cm Hg} - 10.0 \text{ cm Hg} = 9.2 \text{ mm Hg}$
 $HCl \text{ Height (mm)} \times 0.0772$
 $= 7.1 \text{ mm Hg}$

Density Hg is 12.95 times > density HCl(aq)

0.772 mm Hg/cm of acid solution

Density Hg is 12.95 times > density HCl(aq)

$P_{H_2(g)} = 761.5 \text{ mm Hg (barometric)}$
 $- 19.8 \text{ mm Hg } H_2O(g) - 7.1 \text{ mm Hg } HCl(g)$
 $= 734.6 \text{ mm Hg}$
 $= 734.6 \text{ mm Hg} / 760.0 \text{ mm Hg} / 1.000 \text{ atm}$
 $= 0.9666 \text{ atm}$



Moles : Ideal Gas Law
(Part I: Hydrogen Calculation)

$Zn(s) + 2HCl(aq) \rightarrow ZnCl_2(aq) + H_2(g)$

$n_{H_2(g)} = PV / RT$

- $n = \text{moles } H_2(g)$
- $P_{H_2(g)} = 0.9666 \text{ atm}$
- $V = 0.0815 \text{ L}$
- $T = 295.1 \text{ K}$

$R = 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$

$n_{H_2(g)} = 0.00325 \text{ moles } H_2(g) = 0.00325 \text{ moles } Zn(s)$

% Error
Theoretical Mass Zinc vs. Experimental
(Part I: Calculation)

$Zn(s) + 2HCl(aq) \rightarrow ZnCl_2(aq) + H_2(g)$

$\text{mass (g) } Zn(s) = \text{mol } Zn(s) \times \text{Molar Mass } Zn(s)$
 $= 0.00325 \text{ moles } Zn(s) \times 65.37 \text{ g/mol } Zn(s)$

$\% \text{ Error} = \frac{\text{experimental grams } Zn(s) - \text{theoretical grams } Zn(s)}{\text{theoretical grams } Zn(s)} \times 100$
 $= \frac{0.213 \text{ g} - 0.21 \text{ g}}{0.21 \text{ g}} \times 100$
 $= 1.4 \%$


Bring completed Report Forms to Dr. R. to get Mg(s) sample(s).

Molar Mass of any Gas
(Hydrogen for example)

- $PV = nRT$
- $n = g \text{ of gas} / MM_{\text{gas}} [MM_{\text{gas}} = g/mol]$
- $PV = (g \text{ of gas} / MM_{\text{gas}})RT$
- $MM_{\text{gas}} = g \text{ of gas} / V (RT/P)$

Density of gas

- $MM_{\text{gas}} = g \text{ of gas} / V (RT/P)$
- $MM_{\text{gas}} = \text{density of gas} (RT/P)$



(Part II) Magnesium

$Mg(s) + 2HCl(aq) \rightarrow MgCl_2(aq) + H_2(g)$

Mole Calculations:

- Stoichiometry Calculation
- Ideal Gas Law Calculations
- Comparison (% Error)

Get equipment from stockroom and complete data acquisition for Part II.

Have individual Report Forms checked before leaving lab today.

Form II - Mass of Magnesium

Student Name: _____
Section: _____

DATA COLLECTED	
Unknown number	
Volume of hydrogen collected*	mL
Temperature of hydrogen*	°C
Barometric pressure*	mm Hg
Height of solution in eudiometer from benchtop	cm
Height of solution in beaker from benchtop	cm
CALCULATIONS AND RESULTS	
Difference in liquid levels at solution in eudiometer and beaker*	cm Acid Solution
Hydrogen vapor pressure at temperature of hydrogen	mm Hg
Pressure exerted by acid solution*	mm Hg
Difference in mm* (0.772 mm Hg/cm)	mm Hg
Pressure of hydrogen gas*	mm Hg
Mass of hydrogen*	moles
Mass of magnesium*	moles
Mass of magnesium*	g

Show the calculations for each of the entries in the Data Table marked with * on the calculations page.