

Solutions I

Concentrations


Dr. Ron Rusay




Except where otherwise [noted](#), content on this site is licensed under a [Creative Commons Attribution 4.0 International license](#).

Solutions

- ❁ Homogeneous solutions are comprised of **solute**(s), the substance(s) dissolved, [The lesser amount of the component(s) in the mixture], and
- ❁ **solvent**, the substance present in the largest amount.
- ❁ Solutions with less solute dissolved than is physically possible are referred to as “**unsaturated**”. Those with a maximum amount of solute are “**saturated**”.
- ❁ Occasionally there are extraordinary solutions that are “**supersaturated**” with more solute than normal



Concentration and Temperature



Relative Solution Concentrations:

- Saturated**
- Unsaturated**
- Supersaturated**

Dilute


Concentrated

A solution of 35g of potassium chloride in 100g H₂O @ 25°C is Saturated & Concentrated; @ 75°C it is Unsaturated but Concentrated.

What describes
a solution of
25.0g NaCl in
0.100L of H₂O
@ 10°C?

QUESTION


- A) Dilute
- B) Concentrated
- C) Saturated
- D) Unsaturated



What describes
a solution of
100.0g sucrose
in 0.100L of
 H_2O @ 10°C ?


QUESTION

- A) Dilute
- B) Concentrated
- C) Saturated
- D) Unsaturated




DHMO, dihydromonoxide : *“The Universal” Solvent*

<http://www.dhmo.org>



Water : “The Universal” Solvent


The oil (nonpolar) and water (polar) mixture don't mix and are **immiscible**. If liquids form a homogeneous mixture, they are **miscible**.



General
nonpolar
oil and

and
fig.

Water : “The Universal” Solvent



Generally, likes dissolve likes, i.e. polar (water)-polar (solutes) “*hydrophilic*”

Water (polar) repels nonpolar solutes “*hydrophobic*”

QUESTION

An unknown substance dissolves readily in water but not in benzene (a nonpolar solvent). Molecules of what type are present in the substance?


- a) neither polar nor nonpolar
- b) polar
- c) either polar or nonpolar
- d) nonpolar
- e) none of these

Aqueous Reactions & Solutions


- ❁ *Many reactions are done in a homogeneous liquid or gas phase which generally improves reaction rates.*
- ❁ *The prime medium for many inorganic reactions is water which serves as a solvent (the substance present in the larger amount), but does not react itself.*
- ❁ *The substance(s) dissolved in the solvent is (are) the solute(s). Together they comprise a solution. The reactants would be the solutes.*
- ❁ *Reaction solutions typically have less solute dissolved than is possible and are “unsaturated”.*



Salt dissolving in a glass of water



Water dissolving an ionic solid



How can a salt solute be separated from a solution?

<http://chemconnections.org/crystals/>

Solution Concentrations

$$\text{molarity} = M = \frac{\text{moles solute}}{\text{liters solution}}$$

$$\% \text{ by mass} = \frac{\text{mass solute}}{\text{mass solution}} \times 100$$

$$\% \text{ by volume} = \frac{\text{volume solute}}{\text{volume solution}} \times 100$$

[Proof = % by volume x 2]

$$\text{parts per million} = \text{ppm} = \frac{\text{mass solute}}{\text{mass solution}} \times 10^6$$

$$\text{parts per billion} = \text{ppb} = \frac{\text{mass solute}}{\text{mass solution}} \times 10^9$$

$$\text{molality} = m = \frac{\text{moles solute}}{\text{kilograms solvent}}$$




Solution Concentrations

$$\text{Molarity (M)} = \text{moles solute} / \text{Liter}_{\text{solution}}$$

- ❁ *What is the molarity of a solution of 0.50 mol NiCl_2 in 200.0 mL of solution?*

$$M_{\text{NiCl}_2} = [0.50 \text{ mol}_{\text{NiCl}_2} / 200.0 \cancel{\text{mL}}] \times [1000 \cancel{\text{mL}} / \text{L}]$$

$$= 0.25 \text{ mol}_{\text{NiCl}_2} / \text{L}$$




Solution Concentrations

$$\text{Molarity (M)} = \text{moles solute} / \text{Liter}_{\text{solution}}$$

✿ *What is the molarity of a solution of 10.00 g NiCl_2 in 200.0 mL of solution?*

$$M_{\text{NiCl}_2} = [10.00\text{g}_{\text{NiCl}_2} / 200.0\text{mL}] [1\text{mol}_{\text{NiCl}_2} / 129.5\text{g}_{\text{NiCl}_2}] [1000\text{mL} / \text{L}]$$

$$= 0.386 \text{ mol}_{\text{NiCl}_2} / \text{L}$$



Solute Amount
(moles)

1.0



0.079 mol

0


Solution Volume
(Liters)

1.0



0.200 L

0.2



Solution Concentration
(Molarity)

5.0




0.395 M

0

☒ Show values

Solute:

☒ Nickel(II) chloride



QUESTION

40.0-g of HF [MM = 20.0 g/mol] was dissolved in water to give 2.0×10^2 mL of HF(aq), a weak acid solution. The concentration of the solution is:

- a) 0.5 M
- b) 1.0 M
- c) 2.0 M
- d) 5.0 M
- e) 10. M


Solution Concentrations

- ✿ *Mass percent, eg. Glucose, Saline and Ringer's lactate solutions*

$$\text{Mass \%} = \text{Mass solute} / [\text{Mass solute} + \text{Mass solvent}] \times 100$$

- ✿ *What is the mass % of 65.0 g of glucose dissolved in 135 g of water?*

$$\begin{aligned}\text{Mass \%} &= 65.0 \text{ g} / [65.0 + 135] \text{g} \times 100 \\ &= 32.5 \%\end{aligned}$$



Solution Concentrations

Molarity (M) = moles solute / Liter (Solution)

Ringer's lactate solution


$\text{Na}^{1+} = 0.130 \text{ mol/ L}$

$\text{Cl}^{1-} = 0.109 \text{ mol/ L}$

Lactate anion ($\text{C}_3\text{H}_5\text{O}_3$) $^{1-} = 0.028 \text{ mol/ L}$

$\text{K}^{1+} = 0.004 \text{ mol / L}$

$\text{Ca}^{2+} = 0.0015 \text{ mol/ L}$



After blood loss due to trauma, surgery, or burn, it is used for fluid resuscitation usually @ rate equal to 20 to 30 ml/kg body weight/hour.



Solution Concentrations

Molarity (M) = moles solute / Liter_{solution}

mol = (mol solute / Liter_{solution}) x **Liter**_{solution}

❁ An important relationship is **$M \times V_{\text{solution}} = \text{mol}$**

It is used directly in mass calculations of chemical reactions and in the dilutions of solutions.



QUESTION

Solutions: molarity & volume → mass

How many grams of NaCl are contained in 350. mL of a 0.250 M solution of sodium chloride?

- A) 41.7 g
- B) 5.11 g
- C) 14.6 g
- D) 87.5 g
- E) None of these

ANSWER

B) 5.11 g


Seven Solutions Post Lab Questions

<http://chemconnections.org/general/chem120/solutions-mixes.108.html>

Volume (L) times concentration (mol/L) gives moles. Moles are then converted to grams.


Solution Concentrations:

Solute vs. Ion Concentrations



HCl

1.0M 100% Ionized



Acetic Acid ($HC_2H_3O_2$) < 100% Ionized



Preparation of Solutions

Solution Formation
from a Solid

Solutions II

(Solutions/ Molarity)


Applications / Calculations

Dr. Ron Rusay




Except where otherwise [noted](#), content on this site is licensed under a [Creative Commons Attribution 4.0 International license](#).

Preparing a Standard Solution of a Targeted Molarity, M (mol/L)



Preparation of Solutions used in chemistry



Solution Formation
from a Solid

Molarity (M) = Moles solute / Liter (Solution)

QUESTION

A 51.24-g sample of $\text{Ba}(\text{OH})_2$ [MM= 171.3 g/mol] is dissolved in enough water to make 1.20 liters of solution. What is the molarity of the solution?


- a) 0.300 M
- b) 3.33 M
- c) 0.278 M
- d) 2.49×10^{-1} mol/L
- e) 42.7 g/mL

Solution Concentration

- ❖ *The following formula can be used in dilution calculations:*

$$\mathbf{M_1V_1 = M_2V_2}$$

- ❖ *A concentrated stock solution is much easier to prepare and then dilute rather than preparing a dilute solution directly. Concentrated sulfuric acid is 18.0M. What volume would be needed to prepare 250.mL of a 1.50M solution?*
- ❖ $\mathbf{V_1 = M_2V_2 / M_1}$
- ❖ $\mathbf{V_1 = 1.50\ M \times 250.\ mL / 18.0\ M}$
- ❖ $\mathbf{V_1 = 20.8\ mL}$



QUESTION

What volume of 18.0 M sulfuric acid must be used to prepare 15.5 L of 0.195 M H_2SO_4 ?



- A) 168 mL
- B) 0.336 L
- C) 92.3 mL
- D) 226 mL
- E) None of these

Solution Dilution

Solution Formation
by Dilution

Solution Applications

Scoville Units / Capsaicin



Capsaicin: $C_{18}H_{27}NO_3$


Carolina Reaper

“Thunderclap Headache”

http://en.wikipedia.org/wiki/Scoville_scale

<https://www.youtube.com/watch?v=hrF3jVppfr4>

Reversible Cerebral Vasoconstriction Syndrome (RVCS)



<https://www.youtube.com/watch?v=hrF3jVppfr4>

QUESTION

What happens to the number of moles of $\text{C}_{12}\text{H}_{22}\text{O}_{11}$ (sucrose) when 100.0 mL of a 0.20 M solution is diluted to a final concentration of 0.10 M?

- A) The number of moles of $\text{C}_{12}\text{H}_{22}\text{O}_{11}$ decreases.**
- B) The number of moles of $\text{C}_{12}\text{H}_{22}\text{O}_{11}$ increases.**
- C) The number of moles of $\text{C}_{12}\text{H}_{22}\text{O}_{11}$ does not change.**
- D) There is insufficient information to answer the question.**

Solution Applications

A solution of barium chloride was prepared by dissolving 26.0287 g in water to make 500.00 mL of solution. What is the concentration of the barium chloride solution? $M_{\text{BaCl}_2} = ?$

$$M_{\text{BaCl}_2} =$$

$$= [26.0287\text{g}_{\text{BaCl}_2} / 500.00\text{mL}] [1\text{mol}_{\text{BaCl}_2} / 208.23\text{g}_{\text{BaCl}_2}] [1000\text{mL} / \text{L}]$$

$$= 0.25000 \text{ mol} / \text{L}$$



Solution Applications

10.00 mL of this solution was diluted to make exactly 250.00 mL of solution which was then used to react with a solution of potassium sulfate. What is the concentration of the diluted solution. $M_2 = ?$

$$M_{\text{BaCl}_2} = M_1$$

$$M_2 = M_1 V_1 / V_2$$

$$M_2 = 0.25000 \text{ M} \times 10.00 \text{ mL} / 250.00 \text{ mL}$$

$$M_2 = 0.010000 \text{ M}$$



QUESTION

A 51.24-g sample of $\text{Ba}(\text{OH})_2$ is dissolved in enough water to make 1.20 liters of solution. How many mL of this solution must be diluted with water in order to make 1.00 liter of 0.100 molar $\text{Ba}(\text{OH})_2$?

- a) 400. mL
- b) 333 mL
- c) 278 mL
- d) 1.20×10^3 mL
- e) 285 mL

(Chem 120 Prep)

Solution Applications

20.00 mL of a $M_2 = 0.010000$ M barium chloride solution required 15.50 mL of the potassium sulfate solution to react completely. $M_{K_2SO_4} = ?$



$$?M_{K_2SO_4} = [M_{BaCl_2} \times V_{BaCl_2} / V_{K_2SO_4}] [? \text{ mol}_{K_2SO_4} / ? \text{ mol}_{BaCl_2}]$$


$$?M_{K_2SO_4} = \frac{0.010000 \text{ mol}_{BaCl_2} \times 0.02000 \text{ L}_{BaCl_2} \times 1 \text{ mol}_{K_2SO_4}}{L_{BaCl_2} \times 0.01550 \text{ L}_{K_2SO_4} \times 1 \text{ mol}_{BaCl_2}}$$

$$?M_{K_2SO_4} = 0.01290 \text{ mol}_{K_2SO_4} / L_{K_2SO_4} = 0.01290 M_{K_2SO_4}$$



Solution Applications

How many grams of potassium chloride are produced?



$$? \text{ g}_{\text{KCl}} = 0.010000 \text{ mol}_{\text{BaCl}_2} / \text{L}_{\text{BaCl}_2} \times 0.02000 \text{ L}_{\text{BaCl}_2} \times 2 \text{ mol}_{\text{KCl}} / 1 \text{ mol}_{\text{BaCl}_2} \times 74.55 \text{ g}_{\text{KCl}} / \text{mol}_{\text{KCl}}$$


$$= 0.02982 \text{ g}_{\text{KCl}}$$



(Chem 120 Prep)

Solution Applications

If **20.00** mL of a **0.10** M solution of barium chloride was reacted with **15.00** mL of a **0.20** M solution of potassium sulfate, what would be the theoretical yield of barium sulfate?



Which is the Limiting Reagent?

$$\begin{aligned}\text{Mol}_{\text{BaCl}_2} &= M_{\text{BaCl}_2} \times V_{\text{BaCl}_2} \\ &= \frac{0.10 \text{ mol}_{\text{BaCl}_2} / \text{L}_{\text{BaCl}_2} \times 0.02000 \text{ L}_{\text{BaCl}_2}}{1 \text{ mol}_{\text{BaCl}_2}}\end{aligned}$$

$$= 2.0 \times 10^{-3}$$

$$\begin{aligned}\text{Mol}_{\text{K}_2\text{SO}_4} &= M_{\text{K}_2\text{SO}_4} \times V_{\text{K}_2\text{SO}_4} \\ &= \frac{0.20 \text{ mol}_{\text{K}_2\text{SO}_4} / \text{L}_{\text{K}_2\text{SO}_4} \times 0.01500 \text{ L}_{\text{K}_2\text{SO}_4}}{1 \text{ mol}_{\text{K}_2\text{SO}_4}}\end{aligned}$$

$$= 3.0 \times 10^{-3}$$

$$2.0 \times 10^{-3} < 3.0 \times 10^{-3}$$

2.0 x 10⁻³ mol is limiting



(Chem 120 Prep)

Solution Applications

If **20.00** mL of a **0.10** M solution of barium chloride was reacted with **15.00** mL of a **0.20** M solution of potassium sulfate, what would be the theoretical yield of barium sulfate?



Must use the limiting reagent:

$$= \frac{0.10 \cancel{\text{mol}}_{\text{BaCl}_2} \times 0.02000 \cancel{\text{L}}_{\text{BaCl}_2} \times 1 \cancel{\text{mol}}_{\text{BaSO}_4} \times 233.39 \text{ g}_{\text{BaSO}_4}}{1 \cancel{\text{mol}}_{\text{BaCl}_2} \times 1 \cancel{\text{mol}}_{\text{BaSO}_4}}$$

$$= 0.47 \text{ g}$$



QUESTION

What mass of NaOH is required to react exactly with 25.0 mL of 1.2 M H_2SO_4 ?

- A) 1.2 g
- B) 1.8 g
- C) 2.4 g
- D) 3.5 g
- E) None of these

ANSWER

C) 2.4 g

-

Remember that the reaction is $2\text{NaOH} + \text{H}_2\text{SO}_4 \rightarrow \text{Na}_2\text{SO}_4 + 2\text{H}_2\text{O}$, so there are two moles of NaOH used per one mole of H_2SO_4 .