

Acid-Base Equilibrium

Dr. Ron Rusay

© Copyright 2003-2010 R.J. Rusay

Introduction to Aqueous Acids

- ▶ *Acids: taste sour and cause certain dyes to change color.*

Introduction to
Aqueous Acids


Introduction to Aqueous Bases

- ▶ *Bases: taste bitter, feel soapy and cause certain dyes to turn color.*

Introduction to
Aqueous Bases


Models of Acids and Bases

- ▶ *Arrhenius: Acids produce H^+ & bases produce OH^- ion in aqueous solutions .*
- ▶ *Brønsted-Lowry: Acids are H^+ donors & bases are proton acceptors.*




Lewis Acids and Bases

- ▶ **Lewis Acid:** *electron pair acceptor*
- ▶ **Lewis Base:** *electron pair donor*
- ▶ **Example:**




Lewis Acids and Bases

Lewis Acid-Base Theory



Conjugate Acid/Base Pairs




► *conjugate acid*: formed when the proton is transferred to the base.

► *conjugate base*: everything that remains of the acid molecule after a proton is lost.


Click to reveal the conjugate acid and the conjugate base of each species.

Conjugate acid		Conjugate base
<input type="text"/>	NH_3	<input type="text"/>
<input type="text"/>	H_2O	<input type="text"/>
<input type="text"/>	H_2PO_4^-	<input type="text"/>
<input type="text"/>	OH^-	<input type="text"/>

Strong & Weak Acids: Dissociation Constant (K_a)



$$K_a = \frac{[\text{H}_3\text{O}^+][\text{A}^-]}{[\text{HA}]} = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$$



Acid Strength

Strong Acid:

► Equilibrium position lies far to the right. (HNO_3); $K_a \gg 1$

► Produces a conjugate base. (NO_3^-) and a conjugate acid which are weaker than the starting acid and base (H_2O).

Acid Strength

Strong Acids:

Strong Acid	Formula
Hydrochloric	HCl
Hydrobromic	HBr
Hydroiodic	HI
Nitric	HNO_3
Chloric	HClO_3
Perchloric	HClO_4
Sulfuric *	H_2SO_4

Acid Strength (continued)


Weak Acid:

- Equilibrium lies far to the left. (CH_3COOH); $K_a < 1$
- Yields a stronger (relatively strong) conjugate base than water. (CH_3COO^-)

	ACID	BASE	
100% ionized in H_2O	HCl H_2SO_4 HNO_3	Cl^- HSO_4^- NO_3^-	Negligible
	$\text{H}_3\text{O}^+(\text{aq})$	H_2O	
	HSO_4^- H_3PO_4 HF $\text{HC}_2\text{H}_3\text{O}_2$ H_2CO_3 H_2S H_2PO_4^- NH_4^+ HCO_3^- HPO_4^{2-}	SO_4^{2-} H_2PO_4^- F^- $\text{C}_2\text{H}_3\text{O}_2^-$ HCO_3^- HS^- HPO_4^{2-} NH_3 CO_3^{2-} PO_4^{3-}	
Acid strength increases			Base strength increases
	H_2O OH^- H_2 CH_4	OH^- O^{2-} H^- CH_3^-	100% protonated in H_2O
Negligible			Strong

Weak Acids

- Weak acids are only partially ionized in solution.



$$K_a = \frac{[\text{H}_3\text{O}^+][\text{A}^-]}{[\text{HA}]} \quad \text{or} \quad K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$$

- K_a is the acid dissociation constant.

Percent Ionization

- Percent ionization is a way to assess relative acid strengths.


- For the reaction



$$\% \text{ ionization} = \frac{[\text{H}^+]_{\text{eqm}}}{[\text{HA}]_0} \times 100$$

- Percent ionization relates the $\text{H}_3\text{O}^+(\text{aq})$ equilibrium concentration, $[\text{H}^+]_{\text{eqm}}$, to the initial $\text{HA}(\text{aq})$ concentration, $[\text{HA}]_0$.


The Extent of Dissociation for Strong and Weak Acids



Weak Acids

- The higher percent ionization, the stronger the acid.
- Percent ionization of a weak acid decreases as the molarity of the solution increases.
- For acetic acid, 0.05 M solution is 2.0 % ionized whereas a 0.15 M solution is 1.0 % ionized.

Weak Acids Percent Ionization



QUESTION



Nitric acid, HNO_3 , is considered to be a strong acid whereas nitrous acid, HNO_2 , is considered to be a weak acid. Which of the statements here is fully correct?

- A. Nitric acid has an aqueous equilibrium that lies far to the right and NO_3^- is considered a weak conjugate base.
- B. Nitric acid has a stronger conjugate base than nitrous acid.
- C. The dissociation of nitrous acid compared to an equal concentration of nitric acid produces more H^+ .
- D. The equilibrium of nitrous acid lies far to the left and the conjugate base is weaker than the conjugate base of nitric acid.

		ACID	BASE		
Acid strength increases ↑	Strong	HCl	Cl^-	Base strength increases ↓	Negligible
		H_2SO_4	HSO_4^-		
		HNO_3	NO_3^-		
		$\text{H}_3\text{O}^+ (\text{aq})$	H_2O		
		HSO_4^-	SO_4^{2-}		
		H_3PO_4	H_2PO_4^-		
		HF	F^-		
		$\text{HC}_2\text{H}_3\text{O}_2$	$\text{C}_2\text{H}_3\text{O}_2^-$		
		H_2CO_3	HCO_3^{2-}		
	Weak	H_2S	HS^-	Weak	
Negligible		H_2PO_4^-	HPO_4^{2-}		
		NH_4^+	NH_3		
		HCO_3^-	CO_3^{2-}		
		HPO_4^{2-}	PO_4^{3-}		
		H_2O	OH^-		
		OH^-	O^{2-}		
		H_2	H^-		
		CH_4	CH_3^-		
					Strong
					100% protonated in H_2O

Bases

- ▶ “Strong” and “weak” are used in the same sense for bases as for acids.
- ▶ Strong = complete dissociation, $K_b \gg 1$ (concentration of hydroxide ion in solution)



Bases (continued)

- ▶ Weak bases have very little dissociation, $K_b < 1$ (little ionization with water)
- ▶ $\text{CH}_3\text{NH}_2(\text{aq}) + \text{H}_2\text{O}(l) \rightleftharpoons \text{CH}_3\text{NH}_3^+(\text{aq}) + \text{OH}^-(\text{aq})$
- ▶ How conductive is $\text{NaOH}(\text{aq})$ vs morphine, $\text{C}_{17}\text{H}_{19}\text{NO}_3(\text{aq})$?

QUESTION

Aniline, $\text{C}_6\text{H}_5\text{NH}_2$, was isolated in the 1800s and began immediate use in the dye industry. What is the formula of the conjugate acid of this base?

- A. $\text{C}_6\text{H}_5\text{NH}_2^+$
- B. $\text{C}_6\text{H}_5\text{NH}_3^+$
- C. $\text{C}_6\text{H}_5\text{NH}^-$
- D. $\text{C}_6\text{H}_5\text{NH}^+$

Acid-Base Strengths

Strong Acid:

Strong Base:

Acid-Base Strengths

Weak Acid:

Weak Base:


Water as an Acid and a Base Self-ionization



Water as an Acid and a Base




▶ Water is amphoteric (it can behave either as an acid or a base).



$$\text{K}_w = 1 \times 10^{-14} \text{ at } 25^\circ\text{C}$$

Water as an Acid and a Base Self-ionization



The pH Scale


- ▶ $\text{pH} \approx -\log[\text{H}^+] \approx -\log[\text{H}_3\text{O}^+]$
- ▶ pH in water ranges from 0 to 14.

$$\text{K}_w = 1.00 \times 10^{-14} = [\text{H}^+][\text{OH}^-]$$

$$\text{pK}_w = 14.00 = \text{pH} + \text{pOH}$$
- ▶ As pH rises, pOH falls (sum = 14.00).
- ▶ There are no theoretical limits on the values of pH or pOH. (e.g. pH of 2.0 M HCl is -0.301)

The Relations Among $[H_3O^+]$, pH, $[OH^-]$, and pOH


	$[H_3O^+]$	pH	$[OH^-]$	pOH
BASIC	1.0×10^{-15}	15.00	1.0×10^1	-1.00
	1.0×10^{-14}	14.00	1.0×10^0	0.00
	1.0×10^{-13}	13.00	1.0×10^{-1}	1.00
	1.0×10^{-12}	12.00	1.0×10^{-2}	2.00
	1.0×10^{-11}	11.00	1.0×10^{-3}	3.00
	1.0×10^{-10}	10.00	1.0×10^{-4}	4.00
NEUTRAL	1.0×10^{-9}	9.00	1.0×10^{-5}	5.00
	1.0×10^{-8}	8.00	1.0×10^{-6}	6.00
	1.0×10^{-7}	7.00	1.0×10^{-7}	7.00
	1.0×10^{-6}	6.00	1.0×10^{-8}	8.00
ACIDIC	1.0×10^{-5}	5.00	1.0×10^{-9}	9.00
	1.0×10^{-4}	4.00	1.0×10^{-10}	10.00
	1.0×10^{-3}	3.00	1.0×10^{-11}	11.00
	1.0×10^{-2}	2.00	1.0×10^{-12}	12.00
	1.0×10^{-1}	1.00	1.0×10^{-13}	13.00
	1.0×10^0	0.00	1.0×10^{-14}	14.00
	1.0×10^1	-1.00	1.0×10^{-15}	15.00



QUESTION

In a solution of water at a particular temperature the $[H^+]$ may be $1.2 \times 10^{-6} M$. What is the $[OH^-]$ in the same solution? Is the solution acidic, basic, or neutral?

- A. $1.2 \times 10^{-20} M$; acidic
- B. $1.2 \times 10^{-20} M$; basic
- C. $8.3 \times 10^{-9} M$; basic
- D. $8.3 \times 10^{-9} M$; acidic



QUESTION

An environmental chemist obtains a sample of rainwater near a large industrial city. The $[H^+]$ was determined to be $3.5 \times 10^{-6} M$. What is the pH, pOH, and $[OH^-]$ of the solution?

- A. pH = 5.46 ; pOH = 8.54; $[OH^-] = 7.0 \times 10^{-6} M$
- B. pH = 5.46 ; pOH = 8.54; $[OH^-] = 2.9 \times 10^{-9} M$
- C. pH = 12.56 ; pOH = 1.44 ; $[OH^-] = 3.6 \times 10^{-2} M$
- D. pH = 8.54; pOH = 5.46; $[OH^-] = 2.9 \times 10^{-9} M$

The pH Scale

$[H^+]$	$[OH^-]$	pH	pOH	acidic or basic?
$7.5 \times 10^{-3} M$	1.3×10^{-12}	2.1	11.9	Acid
2.8×10^{-5}	$3.6 \times 10^{-10} M$	4.6	9.4	Acid
5.62×10^{-9}	1.78×10^{-6}	8.25	5.75	Base
5.00×10^{-9}	2.00×10^{-6}	8.30	5.70	Base

Indicators

Natural Indicators

Acid-Base Indicators

	pH range for color change													
	0	2	4	6	8	10	12	14						
Methyl violet	Yellow												Violet	
Thymol blue		Red			Yellow					Yellow			Blue	
Methyl orange			Red			Yellow								
Methyl red				Red			Yellow							
Bromthymol blue					Yellow			Blue						
Phenolphthalein							Colorless				Pink			
Alizarin yellow R								Yellow					Red	

QUESTION


Most acid-base indicators are weak acids. In a titration of 0.50 *M* acetic acid (at 25°C, $K_a = 1.8 \times 10^{-5}$) with KOH, which indicator would best indicate the pH at the equivalence point? The approximate K_a for each choice is provided.

- A. Bromophenol blue; $K_a \sim 1 \times 10^{-4}$
- B. Methyl red; $K_a \sim 1 \times 10^{-5}$
- C. Bromothymol blue; $K_a \sim 1 \times 10^{-7}$
- D. Thymolphthalein; $K_a \sim 1 \times 10^{-10}$

Methods for Measuring the pH of an Aqueous Solution



(a) pH paper



(b) Electrodes of a pH meter

QUESTION

The acid-base indicator bromocresol purple has an interesting yellow-to-purple color change. If the approximate K_a of this indicator is 1.0×10^{-6} , what would be the ratio of purple $[A^-]$ to yellow $[HA]$ at a pH of 4.0?

- A. 100:1
- B. 1:100
- C. 1:1
- D. This choice indicates that I don't know.

	ACID	BASE	
100% ionized in H ₂ O	HCl	Cl ⁻	Strong
	H ₂ SO ₄	HSO ₄ ⁻	
	HNO ₃	NO ₃ ⁻	Negligible
	H ₃ O ⁺ (aq)	H ₂ O	
	HSO ₄ ⁻	SO ₄ ²⁻	
	H ₃ PO ₄	H ₂ PO ₄ ⁻	
	HF	F ⁻	
	HC ₂ H ₃ O ₂	C ₂ H ₃ O ₂ ⁻	
	H ₂ CO ₃	HCO ₃ ⁻	
	H ₂ S	HS ⁻	
	H ₂ PO ₄ ⁻	HPO ₄ ²⁻	
	NH ₄ ⁺	NH ₃	
	HCO ₃ ⁻	CO ₃ ²⁻	
	HPO ₄ ²⁻	PO ₄ ³⁻	
	H ₂ O	OH ⁻	
	OH ⁻	O ²⁻	
	H ₂	H ⁻	
	CH ₄	CH ₃ ⁻	
			100% protonated in H ₂ O

The Relations Among $[H_3O^+]$, pH, $[OH^-]$, and pOH

	$[H_3O^+]$	pH	$[OH^-]$	pOH
BASIC	1.0×10^{-15}	15.00	1.0×10^{-1}	-1.00
	1.0×10^{-14}	14.00	1.0×10^{-2}	0.00
	1.0×10^{-13}	13.00	1.0×10^{-3}	1.00
	1.0×10^{-12}	12.00	1.0×10^{-4}	2.00
	1.0×10^{-11}	11.00	1.0×10^{-5}	3.00
	1.0×10^{-10}	10.00	1.0×10^{-6}	4.00
NEUTRAL	1.0×10^{-9}	9.00	1.0×10^{-5}	5.00
	1.0×10^{-8}	8.00	1.0×10^{-6}	6.00
ACIDIC	1.0×10^{-7}	7.00	1.0×10^{-7}	7.00
	1.0×10^{-6}	6.00	1.0×10^{-8}	8.00
	1.0×10^{-5}	5.00	1.0×10^{-9}	9.00
	1.0×10^{-4}	4.00	1.0×10^{-10}	10.00
	1.0×10^{-3}	3.00	1.0×10^{-11}	11.00
	1.0×10^{-2}	2.00	1.0×10^{-12}	12.00
	1.0×10^{-1}	1.00	1.0×10^{-13}	13.00
	1.0×10^0	0.00	1.0×10^{-14}	14.00
	1.0×10^1	-1.00	1.0×10^{-15}	15.00

↑ MORE BASIC
↓ MORE ACIDIC

Conjugates

Some Conjugate Acid-Base Pairs

Acid	K_a	Base	K_b
HNO ₃	(Strong acid)	NO ₃ ⁻	(Negligible basicity)
HF	6.8×10^{-4}	F ⁻	1.5×10^{-11}
HC ₂ H ₃ O ₂	1.8×10^{-5}	C ₂ H ₃ O ₂ ⁻	5.6×10^{-10}
H ₂ CO ₃	4.3×10^{-7}	HCO ₃ ⁻	2.3×10^{-8}
NH ₄ ⁺	5.6×10^{-10}	NH ₃	1.8×10^{-5}
HCO ₃ ⁻	5.6×10^{-11}	CO ₃ ²⁻	1.8×10^{-4}
OH ⁻	(Negligible acidity)	O ²⁻	(Strong base)

$K_a \times K_b = ?$
 $K_a \times K_b = K_w$

Conjugates

$K_a \times K_b = K_w$

HF
 $K_a = 3.5 \times 10^{-4}$

[Click to see conjugate base](#)

What do pK_a and pK_b refer to?
 $pK_a + pK_b = ?$
 $pK_a + pK_b = pK_w$

QUESTION

Values of K_a for Some Common Monoprotic Acids

Formula	Name	Value of K_a
HSO ₄ ⁻	Hydrogen sulfate ion	1.2×10^{-2}
HClO ₄	Chloric acid	1.2×10^{-1}
HC ₂ H ₃ O ₂	Monochloroacetic acid	1.35×10^{-3}
HF	Hydrofluoric acid	7.2×10^{-4}
HNO ₂	Nitrous acid	4.0×10^{-4}
HC ₂ H ₃ O ₂	Acetic acid	1.8×10^{-5}
[Al(H ₂ O) ₆] ³⁺	Hydrated aluminum(III) ion	1.4×10^{-5}
HOCl	Hypochlorous acid	3.5×10^{-8}
HCN	Hydrocyanic acid	6.2×10^{-10}
NH ₄ ⁺	Ammonium ion	5.6×10^{-10}
HOCH ₃	Phenol	1.6×10^{-10}

↑ Increasing acid strength

Use information on this table to determine which of the following bases would have the weakest conjugate acid:

OC₆H₅⁻; C₂H₃O₂⁻; OCl⁻; NH₃

A. OC₆H₅⁻
 B. C₂H₃O₂⁻
 C. OCl⁻
 D. NH₃

Strong vs. Weak Acids

pH Estimations/ Calculations

What are the respective pH values for a 0.100M solution of HCl ($K_a = \infty$) and a 0.100M solution of HF ($K_a = 3.53 \times 10^{-4}$)?

- What are the respective equilibrium concentrations of H⁺ (H₃O⁺)?
- pH is calculated from the equilibrium concentration of H⁺ (H₃O⁺)
- Using K_a , and the starting molarity of acid, the equilibrium concentration of H⁺ (H₃O⁺) can be estimated and then pH; Strong acids 100%, **pH=1.00**, Weak: less than 100%

Strong vs. Weak Acids

pH Estimations/ Calculations

What are the respective pH values for a 0.100M solution of HCl ($K_a = \infty$) and a 0.100M solution of HF ($K_a = 3.53 \times 10^{-4}$)?

- Using K_a , and the starting molarity of the weak acid, the equilibrium concentration of H⁺ (H₃O⁺) can be estimated using an ICE approach and then the pH.

$K_a = [H^+][A^-] / [HA - x] = x^2 / (0.100 M - x)$

$3.53 \times 10^{-4} = x^2 / 0.100$; estimate @ $x \approx (10^{-5})^{1/2}$ representing the [H⁺], taking -log yields a pH >2 and <3.

QUESTION

Which of the following correctly compares strength of acids, pH, and concentrations?


- A. A weak acid, at the same concentration of a strong acid, will have a lower pH.
- B. A weak acid, at the same concentration of a strong acid, will have the same pH.
- C. A weak acid, at a high enough concentration more than a strong acid, could have a lower pH than the strong acid.
- D. A weak acid, at a concentration below a strong acid, could have a lower pH than a strong acid.

Weak Acids K_a and Calculating pH

- ▶ Write the balanced chemical equation clearly showing the equilibrium.
- ▶ Write the equilibrium expression. Use the value for K_a
- ▶ Let $x = [H^+]$; substitute into the equilibrium constant expression and solve.
- ▶ Convert $[H^+]$ to pH.


Equilibrium Concentration Calculations pH from Initial Concentrations and K_a

What is the pH value for a 0.100M solution of HF ($K_a = 3.53 \times 10^{-4}$)?



$$K_a = \frac{[H^+][F^-]}{[HF]}$$

Equilibrium Concentration Calculations



Concentration (M)	HF	H^+	F^-
Initial	0.100	0	0
Change	$0.100 - x$	$+x$	$+x$
Final	$0.100 - x$	x	x

$$K_c = \frac{[H^+][F^-]}{[HF]} = 3.53 \times 10^{-4} = \frac{x^2}{(0.100 - x)}$$

$$3.53 \times 10^{-4} (0.100 - x) = x^2$$

Quadratic:

$$0 = x^2 + 3.53 \times 10^{-4} x - 3.53 \times 10^{-5}$$

$$x = [H^+] = 0.00805 \text{ M}; \text{pH} = 2.09$$

Simplified:

$$3.53 \times 10^{-4} = \frac{x^2}{(0.100)}$$

$$3.53 \times 10^{-4} (0.100) = x^2$$

$$x = [3.53 \times 10^{-4} (0.100)]^{1/2}$$

$$x = [H^+] = 0.00594 \text{ M}; \text{pH} = 2.23$$

QUESTION

Butyric acid is a weak acid that can be found in spoiled butter. The compound has many uses in synthesizing other flavors. The K_a of $HC_4H_7O_2$ at typical room temperatures is 1.5×10^{-5} . What is the pH of a 0.20 M solution of the acid?

- A. 5.52
- B. 4.82
- C. 2.76
- D. -0.70

QUESTION

A 0.35 M solution of an unknown acid is brought into a lab. The pH of the solution is found to be 2.67. From this data, what is the K_a value of the acid?

- A. 6.1×10^{-3}
- B. 1.3×10^{-5}
- C. 7.5×10^{-4}
- D. 2.1×10^{-3}

Behavior of Salts in Water			
The Behavior of Salts in Water			
Salt Solution (Examples)	pH	Nature of Ions	Ion That Reacts with Water
Neutral [NaCl, KBr, Ba(NO ₃) ₂]	7.0	Cation of strong base Anion of strong acid	None
Acidic [NH ₄ Cl, NH ₄ NO ₃ , CH ₃ NH ₃ Br]	<7.0	Cation of weak base Anion of strong acid	Cation
Acidic [Al(NO ₃) ₃ , CrCl ₃ , FeBr ₃]	<7.0	Small, highly charged cation Anion of strong acid	Cation
Basic [CH ₃ COONa, KF, Na ₂ CO ₃]	>7.0	Cation of strong base Anion of weak acid	Anion

LiNO₃

NH₄Cl

Ca(NO₃)₂

MgSO₄

AgClO₃

CsCN


QUESTION

The following salts were all placed in separate solutions at the same temperature so that their concentrations were all equal. Arrange them in order from lowest pH to highest pH.



Additional information: K_b for NH₃ = 1.8×10^{-5} ; K_a for HC₂H₃O₂ = 1.8×10^{-5} ; K_a for Al(H₂O)³⁺ = 1.4×10^{-5} .

- A. NaCl; NH₄NO₃; Ca(C₂H₃O₂)₂; AlCl₃
 B. AlCl₃; NaCl; NH₄NO₃; Ca(C₂H₃O₂)₂
 C. AlCl₃; NH₄NO₃; NaCl; Ca(C₂H₃O₂)₂
 D. NH₄NO₃; AlCl₃; NaCl; Ca(C₂H₃O₂)₂



K_a Values of Some Hydrated Metal Ions at 25°C	
Ion	K_a
Fe ³⁺ (aq)	6×10^{-3}
Sn ²⁺ (aq)	4×10^{-4}
Cr ³⁺ (aq)	1×10^{-4}
Al ³⁺ (aq)	1×10^{-5}
Be ²⁺ (aq)	4×10^{-6}
Cu ²⁺ (aq)	3×10^{-8}
Pb ²⁺ (aq)	3×10^{-8}
Zn ²⁺ (aq)	1×10^{-9}
Co ²⁺ (aq)	2×10^{-10}
Ni ²⁺ (aq)	1×10^{-10}



Structure and Acid-Base Properties

- Two important factors that effect acidity in binary compounds:
 - ⌘ Bond Polarity (smaller e.n. differences favor higher acidities)
 - ⌘ Bond Strength (weak bonds favor higher acidity: more protons [hydronium ions] in solution)
 - ⌘ Select & explain which is the stronger acid: HBr vs. HF.

Oxides





- Acidic Oxides (Acid Anhydrides):
 - ⌘ O-X bond is strong and covalent.
 - $\text{SO}_2, \text{NO}_2, \text{CrO}_3$
- Basic Oxides (Basic Anhydrides):
 - ⌘ O-X bond is ionic.
 - $\text{K}_2\text{O}, \text{CaO}$

Strength of Oxyacids



Strength of Oxyacids

► Name the acids:






Is HBrO_3 stronger or weaker than HClO_3 ?

A) stronger or B) weaker

QUESTION

► Rank 1.0M solutions of HBrO , HIO and HClO in order of increasing acidity.






A) $\text{HBrO} < \text{HIO} < \text{HClO}$ B) $\text{HIO} < \text{HBrO} < \text{HClO}$

C) $\text{HClO} < \text{HBrO} < \text{HIO}$ D) $\text{HIO} < \text{HClO} < \text{HBrO}$

QUESTION

► Rank 1.0M solutions of HBrO , HIO and HClO in order of increasing pH.





A) $\text{HBrO} < \text{HIO} < \text{HClO}$ B) $\text{HIO} < \text{HBrO} < \text{HClO}$

C) $\text{HClO} < \text{HBrO} < \text{HIO}$ D) $\text{HIO} < \text{HClO} < \text{HBrO}$

Strength of Acids

Acid	Formula	K_a (25°C)
Acetic	CH_3COOH	1.8×10^{-5}
Chloroacetic	CH_2ClCOOH	1.4×10^{-3}
Dichloroacetic	CHCl_2COOH	3.3×10^{-2}
Trichloroacetic	CCl_3COOH	2×10^{-1}

Strength of Acids



- 1) Is chloroacetic acid more or less acidic than bromoacetic acid?
2) Will its pKa be higher or lower than bromoacetic acid?

A) 1.more 2.higher B) 1.less 2.lower C) 1.less 2.higher D) 1.more 2.lower

QUESTION

Rank the following acids in order of decreasing acidity.

1) $\text{Br-CH}_2\text{COOH}$, 2) $\text{I-CH}_2\text{COOH}$, 3) CH_3COOH
 $pK_a = 2.69$ $pK_a = 3.12$ $pK_a = 4.75$


A) $1 > 2 > 3$ B) $3 > 2 > 1$ C) $2 > 3 > 1$

QUESTION

Ascorbic acid, also known as vitamin C, has two hydrogen atoms that ionize from the acid. $K_{a1} = 7.9 \times 10^{-5}$; $K_{a2} = 1.6 \times 10^{-12}$. What is the pH, and $\text{C}_6\text{H}_6\text{O}_6^{2-}$ concentration of a 0.10 M solution of $\text{H}_2\text{C}_6\text{H}_6\text{O}_6$?

- A. 2.55; $[\text{C}_6\text{H}_6\text{O}_6^{2-}] = 0.050 \text{ M}$
- B. 2.55; $[\text{C}_6\text{H}_6\text{O}_6^{2-}] = 1.6 \times 10^{-12} \text{ M}$
- C. 1.00; $[\text{C}_6\text{H}_6\text{O}_6^{2-}] = 1.6 \times 10^{-12} \text{ M}$
- D. 5.10; $[\text{C}_6\text{H}_6\text{O}_6^{2-}] = 0.050 \text{ M}$

Neutralization Reactions



Would there be a difference in the reaction of HF versus HCl?

Neutralizations / Titrations



Are there differences in the titration of HF versus HCl? 1) mass wise? 2) pH wise?

- A) 1.NO 2.NO B) 1.YES 2.YES C) 1.YES 2.NO D) 1.NO 2.YES